

Ag 813:457

Bulletin 457

May, 1942

# Tobacco Substation at Windsor Report for 1941

P. J. ANDERSON AND T. R. SWANBACK



Connecticut  
Agricultural Experiment Station  
New Haven



3 9153 01062387 6

# Tobacco Substation at Windsor Report for 1941

P. J. ANDERSON AND T. R. SWANBACK



Connecticut  
Agricultural Experiment Station  
New Haven

## CONTENTS

	PAGE
INTRODUCTION .....	227
THE RELATIVE CROP-PRODUCING CAPACITY OF UREA AND COTTONSEED MEAL.....	229
THE EFFECT OF SOME SOURCES OF PHOSPHORUS ON CIGAR LEAF TOBACCO.....	234
Chemical Analyses .....	237
THE EFFECT OF SOURCE ON MAGNESIUM ABSORPTION BY TOBACCO.....	239
Leaf Content of Magnesium and Calcium .....	241
RESIDUAL EFFECT OF STABLE MANURE .....	242
FURTHER EXPERIMENTS ON STARTER SOLUTIONS .....	243
USE OF CARBON BLACK TO ACTIVATE GROWTH EARLY IN THE SEASON.....	244
FURTHER INFORMATION ON IRRIGATION OF TOBACCO.....	246
LUMARITH AS A SUBSTITUTE FOR GLASS IN THE SEED BED SASH.....	247
TOBACCO DISEASES IN 1941 .....	248
Some Experiments on the Control of Downy Mildew .....	248
Exit Wildfire .....	250
DELAYED FROST INJURY TO TRANSPLANTS IN JUNE.....	250
MOSAIC OR CALICO DISEASE OF TOBACCO.....	253
INSECT INVESTIGATIONS DURING 1941 .....	260
Flea Beetle and Thrips Experiment .....	260
Control Experiments on Open Field Tobacco .....	261
Wireworm Control Experiments .....	262
Insect Abundance During 1941 .....	264

# TOBACCO SUBSTATION AT WINDSOR

## Report for 1941

P. J. ANDERSON AND T. R. SWANBACK

**T**HE Twentieth Annual Report of the Tobacco Substation at Windsor is presented herewith. It is not the object of these annual reports to describe in detail all of the work done during the year on each of many tobacco research projects. Most problems require a continuation through several years before a final answer can be given. In general, it seems best to wait until the conclusion of a project before discussing it in the annual report. On some, however, a progress report is considered necessary. A case in point is the study on phosphorus tests described in this bulletin. On account of world conditions, growers can no longer obtain certain phosphates they have been accustomed to use, and other kinds must be substituted. It may help the grower in choosing a substitute to present here the results of the first two years of a test on various phosphatic materials, even though we have not yet drawn final conclusions.

The growing season of 1941 was on the whole unusually favorable for the production of good tobacco in the Valley. A few local hail and wind storms caused considerable damage but these were confined to small areas. Rainfall was favorable in distribution and amount, as indicated in Table 1, and kept the crop growing steadily. There were no serious leaching rains. The curing season was quite dry, with the result that there was practically no pole rot in this year's crop. Most of the tobacco diseases were at a low ebb and there were no serious insect infestations.

TABLE 1. DISTRIBUTION OF RAINFALL IN INCHES AT THE TOBACCO SUBSTATION, WINDSOR, 1941.

May			June			July			August		
BY TEN-DAY PERIODS											
1-10	11-20	21-31	1-10	11-20	21-30	1-10	11-20	21-31	1-10	11-20	21-31
2.38	.45	.62	3.10	.80	.52	2.12	.45	3.07	.76	1.06	.64
BY MONTHS											
3.45			4.42			5.64			2.46		
AVERAGE FOR PRECEDING 19 YEARS											
3.35			3.79			3.74			4.25		

Of the various forms of direct service offered by the Station, growers take most advantage of the free soil testing service. They have learned to take their own samples. They bring these into the laboratory and, after tests have been made for acidity, nitrogen, potash, phosphorus, calcium, magnesium and sometimes other elements, growers often sit down with the Station agronomist and figure out the best fertilizer formulas for their fields, on the basis of the soil tests. In 1941 over 4,000 samples of soil were brought in and tested before the ground became frozen and ended this work.

A sum of \$1,500 was appropriated for material to build a three-car brick garage. Plans were drawn, contracts let and materials delivered, but work was interrupted when the ground froze in December. This much needed addition to the plant will be finished in the spring.

Table 2 shows the acreage and production of each of the three types of Connecticut Valley tobacco for 1941 and a comparison with 1940 and an average of the previous ten years.<sup>1</sup> This table shows a small increase in acreage of Broadleaf, a smaller decrease in acreage of Havana Seed and a small increase in Shade. Both Shade and Havana Seed are well above the ten-year average. Only Havana Seed shows a large increase in poundage over the ten-year average.

TABLE 2. ACREAGE AND PRODUCTION OF CONNECTICUT VALLEY TYPES IN 1940 AND 1941 COMPARED WITH A TEN-YEAR AVERAGE.

Types	Acres			Production in pounds		
	1941	1940	1930-39	1941	1940	1930-39
Broadleaf	8,300	8,000	8,680	13,284,000	12,326,000	13,373,000
Havana Seed	8,100	8,200	7,690	14,166,000	14,111,000	11,658,000
Shade	6,800	6,400	6,170	6,083,000	5,515,000	6,025,000
Total	23,200	22,600	22,540	33,533,000	31,952,000	31,056,000

The impact of the world war on our tobacco industry was already being felt in 1941 and will surely be greater in 1942. First of all, labor became scarce and high priced because of the location of defense industries in the Valley. These firms paid wages the tobacco farmer could not hope to meet.

Prices of materials, machinery, shade cloth, fertilizers, etc., have been climbing. Certain kinds of fertilizer materials which the tobacco grower especially preferred have about gone off the market. Precipitated bone, the favorite source of phosphorus, cannot be obtained. Other forms of bone are also scarce. A shortage of domestic sources of potash was anticipated but has not developed. There seems to be enough domestic sulfate of potash and cottonhull ashes to take care of the potash requirements. Uramon, which has come to be an important source of nitrogen in tobacco mixtures, can no longer be obtained because the DuPont Company, the sole manufacturer of

<sup>1</sup> From release of New England Crop Reporting Service December 23, 1941.

this material, has been obliged to turn so much of its chemical producing capacity to war chemicals. The effect of dislocations of freight and ocean transport on the other materials cannot be predicted at this time.

Since practically all Connecticut-grown tobacco is used domestically, the loss of foreign markets has no effect on it. The cutting off of Sumatra by war in the Pacific, however, may have a profound effect on our Shade industry. From one-third to one-half of the cigars produced in America are wrapped with Sumatra Leaf. There is now a large supply in this country but, if the war should continue to isolate Sumatra until this supply runs out, other sources of cigar wrappers must be found, and the logical substitute is Connecticut Valley Shade. If manufacturers should all turn to Shade the acreage could be almost doubled.

Detailed or summary reports on some of the projects under way at the Substation are presented in the following pages.

#### THE RELATIVE CROP-PRODUCING CAPACITY OF UREA AND COTTONSEED MEAL

The crop-producing capacity of any nitrogenous fertilizer is related to the nitrate concentration which it can develop and maintain in the soil during a relatively short period, from mid-June until early August. Very little nitrate is needed by the tobacco plants previous to this time and any that is supplied after this period—when the crop is already mature or harvested—is obviously wasted. Measurements of the concentration of nitrates in the soils to which such materials have been applied should, therefore, give the first indication of crop-producing efficiency. Such measurements were made at weekly intervals for four years (1932-35) at the Substation on tobacco plots treated with single nitrogen carriers. The results were published in Conn. Sta. Bul. 386, pp. 552-574. Comparing the cottonseed meal figures with those for urea (Table 13 of that report) we find that the average concentration of nitrate in the soil between June 18 and August 13 was 37 pounds per acre for cottonseed meal and 58 pounds for urea, i. e., cottonseed meal was much less effective than urea in its nitrate-producing capacity during these critical few weeks.

That more nitrate nitrogen is liberated from urea than from cottonseed meal and that the crop is able to utilize more of it is also indicated by lysimeter experiments which have been carried on here since 1929. In the lysimeter tests all nitrate nitrogen which is removed from the soil, either by leaching or by the growing crop during the year, is accurately measured.

Results of these 1940-41 lysimeter studies in nitrogen removal by the crop and by leachate are given for urea and cottonseed meal

in Table 3.<sup>1</sup> This table shows that, starting with the same amount of nitrogen supplied in the fertilizer, a greater percentage was always recovered in the crop and that the total recovery (crop and leaching) was also always higher for urea.

In another series of lysimeter tests, 200 pounds per acre of nitrogen, in urea and cottonseed meal, were added to the soil annually for nine years. This was cropped to tobacco. The average annual

TABLE 3. REMOVAL OF NITRATE NITROGEN BY CROP AND BY LEACHING IN 12 MONTHS, 1940-41. COMPARISON OF UREA AND COTTONSEED MEAL IN LYSIMETERS.

Pounds of nitrogen added in fertilizer	Nitrogen carrier	Pounds of nitrogen recovered		
		In crop	Leached as nitrate	Total
0	None	33	47	80
120	Urea	91	58	149
	Cottonseed meal	65	68	133
160	Urea	100	77	177
	Cottonseed meal	77	73	150
200	Urea	130	90	220
	Cottonseed meal	85	84	169

removal of nitrogen (nitrate nitrogen in the leachate and total nitrogen in the plants) was 165 pounds for urea, 145 pounds for cottonseed meal and 47 pounds where no nitrogen was applied as fertilizer. We may assume that the 98 pounds (145 minus 47) of the nitrogen derived from cottonseed meal was sufficient to raise a normal crop. Since 200 pounds of nitrogen as urea liberated 118 pounds in excess of soil production of available nitrogen, then 166 pounds of nitrogen added in this material could supply the required 98 pounds. Thus, in this test, urea was about 20 percent more efficient than cottonseed meal in providing nitrates for crop use.

On the field plots mentioned in the first paragraph in this section, the crop fertilized with urea was always larger than that on the cottonseed meal plots, the latest published report (Bul. 386, p. 546, of this Station) showing an average annual increase of 100 pounds of cured tobacco to the acre for urea during the six preceding years. The tobacco on the urea plots, however, was inclined to be too dark and heavy for best quality. Since this objectionable characteristic often has been observed on plots which were too heavily fertilized with nitrogen from any source, it was natural to conclude that the same cause was operating here and that the ration of nitrates furnished by urea was too great for obtaining the best quality leaves.

<sup>1</sup> Data on the lysimeter experiments furnished by M. F. Morgan of the Soils Department.



Assuming that this conclusion was correct, it seemed desirable to determine by field tests the relative efficiency of the two materials. If 200 pounds of nitrogen to the acre in cottonseed meal be regarded as the standard amount for growing a normal crop of tobacco, how much nitrogen in urea should be used to produce the same results?

To answer this question a series of 18 one-thirtieth acre plots was laid out in 1937, with six different treatments in triplicate. The only differences in the fertilizer applications to these plots were in the quantity and source of nitrogen, all other fertilizer ingredients being kept constant. The various nitrogen treatments were:

150 pounds N in urea	175 pounds N in cottonseed meal
150 pounds N in cottonseed meal	200 pounds N in urea
175 pounds N in urea	200 pounds N in cottonseed meal

All plots received the same cultural treatment and were grown and harvested in the customary way for Havana Seed tobacco. The cured leaves were graded and weighed in the Station warehouse and the grades computed to a single figure (grade index<sup>1</sup>) from the

TABLE 4. EFFICIENCY OF UREA COMPARED WITH COTTONSEED MEAL. FOUR YEARS' SUMMARY OF YIELDS AND GRADING.

Material	Lbs. N per acre	Acre yield					Grade index				
		1937	1939	1940	1941	Av.	1937	1939	1940	1941	Av.
Urea	150	1706	1803	1506	1894	1843	.302	.373	.294	.405	.365
		2045	1916	1532	2206		.367	.393	.349	.407	
		2073	1940	1444	2038		.366	.392	.336	.397	
Urea	175	2095	1940	1495	2116	1924	.396	.391	.349	.429	.369
		2152	2058	1490	2134		.380	.379	.283	.377	
		2077	2092	1419	2014		.369	.397	.291	.381	
Urea	200	1957	2005	1500	2315	1995	.335	.398	.345	.448	.380
		2273	2046	1688	2346		.380	.373	.355	.396	
		2162	2000	1523	2123		.403	.384	.341	.395	
C. S. M.	150	1849	1875	1510	1916	1803	.326	.376	.359	.364	.355
		2033	1872	1605	2058		.326	.388	.380	.385	
		1790	1831	1430	1864		.342	.347	.355	.319	
C. S. M.	175	2004	1830	1514	1857	1842	.366	.358	.370	.377	.360
		2091	1937	1581	2147		.345	.383	.368	.385	
		1916	1823	1515	1886		.345	.361	.328	.340	
C. S. M.	200	2355	1924	1504	2250	1926	.384	.368	.347	.418	.368
		2227	1891	1582	2178		.380	.391	.321	.408	
		1988	1770	1495	1947		.360	.349	.306	.382	

<sup>1</sup> Grade index is a figure which represents the relative value of a lot of tobacco computed on the percentage weight of each grade of leaves in the lot and the relative values of these grades. Assuming that the light wrapper is the perfect leaf of Havana Seed tobacco, we assign to it a value of 100. The other grades are assigned values of the same proportion of 100 as their market value was to the price of the light wrapper when this system was established. Thus medium wrappers have a value of 60; long seconds (19 inches or more) are 60; short seconds (15 and 17 inches) are 30; long darks are 30; dark stemming (short darks of 15 and 17 inches) are 20; fillers and brokes are 10. It is true that the values of these grades have fluctuated considerably during the 15 years that we have used this system of comparing lots of tobacco but, in order to be able to average results over a period of years, it seems advisable to retain the same system for the present. To obtain the grade index figure the percentage of each grade in a lot of tobacco is multiplied by the relative values noted above and the products are added.

weights and relative values of each grade. The results are presented in Table 4 for the years 1937, 1939, 1940 and 1941. The crop of 1938 was omitted from the calculations because the extreme weather conditions of that year resulted in such poor quality that for the most part it was not worth sorting.

From the averages (bold face type) in this table the following observations may be made:

1. Each increase in quantity of nitrogen in the fertilizer, whether in urea or in cottonseed meal, produced an increase in yield.
2. Each increase in fertilizer nitrogen, regardless of source, improved the grade index.
3. When the quantity of nitrogen was the same, both the yield and index were always higher for urea.




Figure 1. Concentration of nitrate in soils of urea and cottonseed meal plots during season of 1937.

4. With 175 pounds of nitrogen in urea practically the same yield and grade index were obtained as with 200 pounds of nitrogen in cottonseed meal.

5. When the urea nitrogen was reduced to 150 pounds, however, the average yield was considerably below that of 200 pounds in cottonseed meal. The grade index was also somewhat lower. In two

out of the four years, however, 150 pounds in urea was as good as 200 in cottonseed meal.

*The results from four years of this trial indicate, therefore, that urea nitrogen is at least 14 percent more efficient than cottonseed meal nitrogen in its crop-producing capacity.*

During one year of this field test (1937) the nitrate concentration in the soil was determined at weekly intervals. The average of all urea plots, contrasted with the cottonseed meal plots, is graphically presented in Figure 1, where it will be seen that urea consistently accumulated more nitrate than did cottonseed meal. The average concentration of nitrates in all urea plots, from June 14 to August 16, was 73.5 pounds of nitrogen per acre, contrasted with 41.6 for the cottonseed meal plots. This suggests that the better yield on the urea plots was due to the greater supply of nitrates available in the soil during the critical period. It will be observed, however, that the difference in nitrate concentration is greater than the percentage increase in crop production, showing that the crop return is not exactly proportional to the quantity of nitrates in the soil. This may be due to the presence of other growth materials in cottonseed meal or to the fact that the yields here are too close to the upper limit of possible crop response to increased fertilizer; i. e., the law of diminishing returns is operating. A more correct conclusion may be drawn by comparing the "175 pound urea" plots with the "200 pound cottonseed meal" plots. These two gave nearly the same crop response, as already noted. The average concentration of the former during the summer was 53 pounds of nitrate-nitrogen per acre while the latter was 48 pounds. This makes a difference of 5 pounds, which is probably too small to be of significance in these field tests.

**Comparative Grading.** An important question to the handler of tobacco is: what effect does the substitution of urea for cottonseed meal have on the percentage of grades when the leaves are sorted?

The sorting records for the crop of 1941 on these plots are shown in Table 5.

This year is selected because it was the final year of the test and should, therefore, show the maximum results of any cumulative effects from continuous use of each fertilizer. Moreover, it was a favorable growing season and resulted in the most regular crop of the series. The data presented in this table show that, contrary to our previous impression, urea did not increase the percentage of dark grades nor decrease the percentage of the better light grades. For the same quantity of fertilizer nitrogen, therefore, the grade index was uniformly higher for urea. No burn tests were made this year because many previous trials at this Station (see particularly Bul. 299, p. 172) have demonstrated that the fire-holding capacity is not affected by the use of urea. Divergent opinions have been ex-

pressed by buyers, packers and manufacturers as to the effect of urea on the more intangible qualities such as taste, aroma, finish and the like. Since there is no known method of measuring these factors of quality, and, moreover, opinion is so diversified, this question does not appear close to solution in the near future.

TABLE 5. YIELD AND GRADING RECORDS OF UREA EFFICIENCY COMPARED WITH COTTONSEED MEAL PLOTS. CROP OF 1941.

Material <sup>1</sup>	Lbs. N per acre	Plot no.	Yield, lbs. per acre		Percentage of grades								Grade index		Crop index	
			Plot	Av.	L	M	LS	SS	LD	DS	F	B	Plot	Av.		
Urea	150	N70	1894		2	2	39	2	38	2	9	6	.405		.403	824.54
		N70-1	2206	2046	4	2	34	3	42	1	8	6	.407			
		N70-2	2038		2	2	37	3	37	4	12	3	.397			
Urea	175	N71	2116		6	2	37	2	37	2	9	5	.429		.396	826.85
		N71-1	2134	2088	1	1	31	7	46	2	11	1	.377			
		N71-2	2014		1	1	34	4	43	3	10	4	.381			
Urea	200	N72	2315		5	1	43	2	39	1	7	2	.448		.413	933.79
		N72-1	2346	2261	3	2	36	2	37	1	8	11	.396			
		N72-2	2123		1	1	38	3	41	3	11	2	.395			
C. S. M.	150	N73	1916		1	1	32	4	39	4	11	8	.364		.356	692.78
		N73-1	2058	1946	2	1	34	4	40	4	10	5	.385			
		N73-2	1864		..	..	26	2	40	5	12	15	.319			
C. S. M.	175	N74	1857		2	2	33	2	37	6	12	6	.377		.367	720.42
		N74-1	2147	1963	2	3	31	4	43	3	10	4	.385			
		N74-2	1886		..	..	26	3	50	4	13	4	.340			
C. S. M.	200	N75	2250		4	4	35	2	41	1	8	5	.418		.403	856.38
		N75-1	2178	2125	3	2	36	3	41	3	9	3	.408			
		N75-2	1947		1	2	35	3	38	6	11	4	.382			

<sup>1</sup> Relative Crop Values.

200 Urea = 109.0

175 Urea = 96.6

150 Urea = 96.3

200 C. S. M. = 100.0

175 C. S. M. = 84.1

150 C. S. M. = 80.9

## THE EFFECT OF SOME SOURCES OF PHOSPHORUS ON CIGAR LEAF TOBACCO.

T. R. SWANBACK, M. F. MORGAN AND P. J. ANDERSON

In the fertilization of tobacco in the Connecticut Valley growers have been accustomed to use bone phosphates in one form or another. Precipitated bone is probably the most commonly used material. Earlier investigations by Jenkins in 1914 (Conn. Sta. Bul. 180:28-30) and later by Nelson and Anderson (Tobacco Sta. Bul. 5:17-24) showed favorable results with precipitated bone. The greater portion of this material is usually imported from Belgium. Due to present war conditions this source of supply is suspended. Even a second choice of bone phosphates, such as raw bone and steamed bone, soon may not be available in sufficient quantities to meet the demand.

In anticipation of such shortage of bone phosphates, an experiment on field plots at the Windsor Substation was begun with the object of finding whether other phosphates could be substituted.

Included in the materials, in addition to precipitated bone and steamed bone, were concentrated superphosphate (triple superphosphate), calcium metaphosphate, potassium metaphosphate and potassium calcium metaphosphate. These were furnished by the Tennessee Valley Authority. The last mentioned material was added to the tests in 1941, while the others were included in the experiment when it was begun in 1940.

The field allotted to the experiment had been used for vegetable crops for many years previous to 1940 and possessed normal fertility. Therefore, it was not anticipated that there would be symptoms of phosphorus deficiency on plots included in the test, even where the fertilizer contained no special carrier of phosphorus.

The various phosphatic materials, each furnishing 116 pounds  $P_2O_5$  per acre, were added to one common fertilizer formula that included uramon, soybean oil meal and nitrate of soda as carriers of nitrogen and potash in the form of sulfate of potash. These materials were selected because of very low content or almost entire lack of  $P_2O_5$ . Lime, magnesium sulfate (Emjeo) and landplaster were also included for the purpose of establishing acid-basic balance and distributing an exact and uniform amount of plant nutrients among the treatments. All tests were carried out in quadruplicate (one-fortieth acre) plots, except the potassium calcium metaphosphate. For tests on the latter material, the size of the field permitted inclusion of only two plots.

The summer of 1941 was a favorable growing season and uniform growth was observed on the entire field. In mid-season, however, it was noticed that tobacco on the calcium metaphosphate plots had a more luxuriant spread and was taller than that under other treatments.

The following general remarks were made at the time of sorting.

- No P-carrier: short, dark, coarse, not veiny.
- Precipitated bone: good to very good quality, not veiny.
- Triple superphosphate: good, veiny, slightly reddish.
- Steamed bone: good, but a little heavy, slightly veiny.
- Ca-metaphosphate: very good quality, slightly veiny.
- K-metaphosphate: very good quality, slightly veiny.
- K, Ca-metaphosphate: very good but short, not veiny.

The yield and grading records will be found in Table 6.

In examining the yield data it is apparent that, in the second year of the experiment, there was no difficulty in obtaining heavy yields on this particular test field, even with no special phosphorus carrier.

Calcium metaphosphate and triple superphosphate produced about 7 percent higher yields than the check; precipitated bone about 5 percent higher, while steamed bone equalled the check. However,

none of these deviations from the yields of the check plots might be considered significant.

On the other hand, potassium metaphosphate and potassium calcium metaphosphate produced an average of 15 percent lower yields than the control plots. It is of interest to keep in mind that potassium is the carrier of phosphorus in both these materials. Potassium calcium metaphosphate contains only about 4 percent CaO and over 35 percent K<sub>2</sub>O; that is, the same as for potassium metaphosphate.

TABLE 6. YIELD AND GRADING RECORDS OF PHOSPHORUS PLOTS. CROP OF 1941.

Treatment	Plot no.	Yield lbs. per acre		Percentage of grades								Grade index	
		Plot	Av.	L	M	LS	SS	LD	DS	F	B	Plot	Av.
No phosphorus carrier	P11	2503		1	1	38	3	46	1	9	1	.403	
	P11-1	2300		..	2	36	3	47	1	8	3	.391	
	P11-2	2040	2352	2	2	33	4	45	2	8	4	.393	.406
	P11-3	2564		3	4	40	3	41	1	7	1	.436	
Precipitated bone	P12	2490		4	3	31	3	48	1	9	1	.409	
	P12-1	2599		2	2	39	2	44	1	8	2	.416	
	P12-2	2657	2457	4	3	41	2	41	..	7	2	.442	.409
	P12-3	2082		..	..	33	4	47	2	11	3	.369	
Triple super-phosphate	P13	2403		1	2	40	3	43	1	8	2	.412	
	P13-1	2602		2	3	39	3	39	1	7	6	.413	
	P13-2	2694	2517	2	..	38	2	48	1	7	2	.409	.412
	P13-3	2369		..	2	42	4	43	1	7	1	.415	
Steamed bone	P14	2511		3	2	36	2	46	1	9	1	.414	
	P14-1	2183		2	1	41	4	42	1	9	..	.421	
	P14-2	2335	2344	1	2	44	2	38	1	7	5	.420	.418
	P14-3	2348		2	2	40	3	40	1	9	3	.415	
Calcium metaphosphate	P15	2472		5	3	38	2	42	1	7	2	.439	
	P15-1	2556		6	3	35	3	42	1	9	1	.435	
	P15-2	2596	2519	3	1	43	2	42	1	7	1	.436	.446
	P15-3	2450		9	4	40	2	33	3	7	2	.474	
Potassium metaphosphate	P16	1984		3	2	39	4	40	1	10	1	.421	
	P16-1	1809		..	..	30	2	41	3	15	9	.339	
	P16-2	2350	2071	4	6	38	4	39	1	7	1	.443	.396
	P16-3	2141		..	..	35	4	48	1	11	1	.380	
Potassium Ca-metaphosphate	P17	1944		5	3	38	3	39	1	9	2	.435	
	P17-1	1883	1914	3	3	36	3	41	2	10	2	.412	.424

Although reasonable yields are of importance in crop production with respect to leaf tobacco, quality is still more important. Of the materials tested, only the calcium metaphosphate produced tobacco decidedly superior in grade to that of the control plots (average grade index .406, as compared with .446). In addition, it is noteworthy that calcium metaphosphate produced about 12 percent less "darks" than the check.

The other sources of phosphorus produced tobacco of somewhat higher grading than the control (two out of four plots of potassium metaphosphate reached higher indexes than the check). The varia-

tion between treatments, however, is not much higher than the variation in values between plots.

If quantity to a certain extent makes up for quality, the crop index (product of yield figure and corresponding value of grade index) should give a conception of the return values of various treatments.

The following resumé, based on average crop indexes and giving a value of 100 to the control, presents an evaluation of the return from the use of the various sources of phosphorus. They may be referred to as relative crop values.

No phosphorus	100.	
Steamed bone	101.6	
Precipitated bone	105.2	
Triple superphosphate	108.6	
Calcium metaphosphate	117.6	
Potassium metaphosphate	98.0	Av. of two good plots
Potassium metaphosphate	74.5	Av. of two inferior plots
Potassium metaphosphate	85.9	Av. of all four plots
Potassium calcium metaphosphate	85.0	Av. of two plots

For comparison, a two-year average of corresponding values is given below (with the exception of potassium calcium metaphosphate, for which data of only one year values are available).

No phosphorus	100
Potassium metaphosphate	100 (at best)
Steamed bone	101.9
Triple superphosphate	104.6
Precipitated bone	106.6
Calcium metaphosphate	113.6

These two-year averages agree closely with results of the present year, with the exception that the positions of precipitated bone and superphosphate are reversed. If it be recalled that the triple superphosphate had a tendency to produce an undesirable reddish cast of the cured leaf, this *per se* reduces the possible value of the material in cigar leaf production, in particular as regards wrapper tobacco.

Therefore, the rank of the phosphorus sources tested, considering their relative quality, would be as follows:

1. Calcium metaphosphate
2. Precipitated bone
3. Steamed bone
4. Potassium metaphosphate
5. Potassium calcium metaphosphate
6. Triple superphosphate

#### Chemical Analyses

Samples of tobacco grown with the various phosphorus carriers were collected from the replicates to make up one composite sample of long seconds and one of long darks per treatment. They were dried, ground to a fine powder and then analyzed with respect to

content of phosphorus, potassium, calcium and magnesium. The results on phosphorus are shown in Table 7.

It is seen that the phosphoric acid content varies little, about half of one percent. Tobacco fertilized with precipitated bone and calcium metaphosphate contained the least phosphorus, particularly in the "seconds." In considering crop removal of phosphorus, however, we find in the following summary a rather uniform utilization of the phosphorus available in the soil.

TABLE 7. PHOSPHORUS CONTENT OF CURED TOBACCO. AIR DRY BASIS.  
CROP OF 1941.

Source of phosphorus	Grade	Plot no.	Percent $P_2O_5$	Av.
None	LS	P11	0.45	0.50
	LD		0.55	
Precipitated bone	LS	P12	0.41	0.47
	LD		0.53	
Triple superphosphate	LS	P13	0.46	0.52
	LD		0.57	
Steamed bone	LS	P14	0.43	0.49
	LD		0.55	
Calcium metaphosphate	LS	P15	0.38	0.45
	LD		0.52	
Potassium metaphosphate	LS	P16	0.49	0.51
	LD		0.53	
Potassium calcium metaphosphate	LS	P17	0.50	0.51
	LD		0.52	

Average removal of phosphoric acid ( $P_2O_5$ ) by yield of leaves from various phosphorus treatments.<sup>1</sup>

Check	11.76 pounds per acre
Precipitated bone	11.55 pounds per acre
Triple superphosphate	13.09 pounds per acre
Steamed bone	11.49 pounds per acre
Calcium metaphosphate	11.33 pounds per acre
Potassium metaphosphate	10.56 pounds per acre
Potassium calcium metaphosphate	9.76 pounds per acre

The check, precipitated bone, steamed bone and calcium metaphosphate come readily within the range of average phosphorus removal (11.56 pounds per acre), while triple superphosphate removed more, and the two potash phosphates removed less, than the average. From this it is suggested that triple superphosphate alone renders more available phosphoric acid than any of the materials tested.

A detailed report on the analytical results on potassium, calcium and magnesium is not given here since no significant differences due to treatments were apparent. The average potash ( $K_2O$ ) content

<sup>1</sup> Stalks remove about as much  $P_2O_5$  as the leaves, hence total crop removal would be about twice the amounts given.



was  $4.09 \pm 0.206$  percent, calcium (CaO)  $5.75 \pm 0.1$  percent and magnesia (MgO)  $1.70 \pm 0.066$  percent. Analytical results give little or no explanation, in particular to the behavior of calcium metaphosphate which ranked the highest in the present investigation. It is possible that this material has a favorable influence on rendering other nutrients more available, and this influence may be extended to reactions in the plant tissues.

### THE EFFECT OF SOURCE ON MAGNESIUM ABSORPTION BY TOBACCO

T. R. SWANBACK

The role of magnesia and its place in cigar leaf production are rather well established through extensive work at this Station. (See Bulls. 350 and 386). The tobacco grower in the Connecticut Valley who avails himself of the soil testing service keeps informed on the status of magnesia content in his land. Although many of the materials used in tobacco fertilizers contain more or less magnesia, it is often necessary to include a special carrier. Under conditions where it has been advisable to use lime containing magnesia, the question has often been asked whether magnesian limestone (dolomite) or hydrated magnesian lime is more effective in producing the desired results in the same season that materials are applied.

In an attempt to find an answer to this question a field was selected which was low in fertility (soil reaction pH 4.8) and which had not grown tobacco for at least 15 years. The field was divided into four one-twentieth acre plots, one of which did not receive any liming material.

The remaining three plots were treated as follows:

	Pounds per acre of		
	Material	CaO	MgO
1. Hydrated magnesian lime	800	300	200
2. Magnesian limestone	1000	300	200
3. {Landplaster <sup>1</sup> {plus Emjco (Anhydr. MgSO <sub>4</sub> )	1000	300	100 190}

The entire field was fertilized uniformly with an 8-4-8 mixture at the rate of 2,500 pounds per acre. Tobacco was planted crosswise to the lime applications.

Vigorous growth was observed on the three treated plots, while the size of plants on the check plot was noticeably smaller.

While the main purpose of growing the crops was to determine the content of magnesium and possibly calcium in the cured leaves from the various applications, it may be of interest to learn about the

<sup>1</sup> Landplaster and magnesium sulfate were included as a treatment for comparing a neutral supply of CaO and MgO versus the lime materials which offset a part of the soil acidity.

effect on yield and grading. On the sorting table all the tobacco from this field was considered to be of very good quality, with satisfactory texture and color, except for the plot receiving landplaster (calcium sulfate) and magnesium sulfate. Here, the fillers (leaves closest to the ground) were of a blackish color (known to the trade as "black tobacco") and, in general, the tobacco from this treatment was inferior in quality to the control and the two lime treatments. In this connection, it should be pointed out that landplaster is *not recommended* for application to tobacco land that is *too acid*. As has been pointed out earlier, (Bul. 350:477 and Bul. 444:260), landplaster may be employed only on tobacco soils having a satisfactory reaction (pH).

Table 8 lists yield and grading data. In line with field observations, it is found that the control plot produced the lowest yield, while hydrated lime and landplaster produced almost identical yields, about 300 pounds per acre more than the control, and magnesium limestone produced about 480 pounds more than the control.

The grading, as indicated by the grade indexes, again shows that the landplaster treatment falls below the rest in quality.

TABLE 8. YIELD AND GRADING RECORDS OF MAGNESIAN LIME TEST. CROP OF 1941.

Treatment	Yield per acre	Percentage of grades								Index	
		L	M	LS	SS	LD	DS	F	B	Grade	Crop
Control	1978	8	2	35	2	38	3	10	2	.440	870.3
Hydrated Mg lime	2295	10	3	40	1	35	1	9	1	.478	1097.0
Landplaster + Emjeo	2280	2	2	36	1	41	2	12	4	.394	898.3
Mg limestone	2460	11	5	34	1	37	1	9	2	.471	1158.7

In considering the crop index (which is the product of the yield figure and corresponding value of the grade index) it is still more evident that the two magnesian lime materials gave a far better return than the control and landplaster treatment.

From computed relative crop values (crop index of control = 100), the following resumé is obtained:

Control	=	100
Landplaster + Emjeo	=	103.2
Hydrated magnesian lime	=	111.5
Magnesian limestone	=	133.1

Once during the growing season (mid-July) and later, after the crop had been removed (October) samples of soil were collected from the four plots. Reaction and content of replaceable calcium and magnesium (Morgan method) were determined on these samples.

The results are found in Table 9. With the exception of magnesian limestone, no appreciable change in reaction was effected

through the use of various liming materials. Replaceable calcium increased during the season, apparently through nitrification or other biological processes in the soil.

If it be assumed that 50 pounds of magnesium per acre was the original content of the soil and that a small amount was incidentally supplied in the fertilizer, all the magnesium supplied as magnesium sulfate (Emjeo) and magnesium limestone was found to be replaceable, while hydrated lime was not so effective in this respect.

TABLE 9. SOIL REACTION AND CONTENT OF REPLACEABLE CALCIUM AND MAGNESIUM IN SOIL OF LIME EXPERIMENT PLOTS.

Treatment	pH		Calcium		Magnesium	
	July	Oct.	July	Oct.	July	Oct.
Control	4.90	5.10	400	800	50	25
Landplaster+Emjeo	4.84	5.10	750	1000	250	100
Hydrated lime	5.04	5.10	750	1000	100	50
Magnesian limestone	5.10	5.50	800	900	250	250

#### Leaf Content of Magnesium and Calcium

Representative samples of lights (L) mediums (M) long seconds (LS) and long darks (LD) from the four treatments were ground up finely and analyzed<sup>1</sup> for calcium and magnesium. The results are given in Table 10. In examining the data it is seen that, on the average, slightly more magnesium was deposited in the leaves when hydrated lime was the source than when the limestone was used.

It is of interest to note that calcium in hydrated lime was less readily taken up by tobacco than that from limestone. This, however, does not necessarily mean that the calcium in limestone is more readily available, but that the magnesia in this material may not have been so readily available as that in the hydrated lime. The reciprocal or antagonistic actions of the three main bases, — calcium, potassium and magnesium — function to establish an acid-basic balance within the plant. That is, any one of the three bases makes up for the insufficiency of one or both of the others (See Bul. 444:256).

Emjeo (magnesium sulfate) and landplaster in the present comparison were responsible for the highest leaf contents of both magnesium and calcium. This is in line with an earlier report (Bul. 386:584) that Emjeo furnishes more readily available magnesia than does magnesian lime. Calcium in landplaster (calcium sulfate), because of its chemical composition, as would be expected furnished more readily available calcium than either of the lime materials.

The results of this investigation suggest that either one of the two materials, magnesian limestone and hydrated lime, may be used

<sup>1</sup> By the Chemistry Department of the New Haven Station.

as a source of magnesia with about equal advantage, although the limestone might be safer to use in fertilizer mixtures of any combination.

It may be concluded that, while there may be a choice between the two lime materials when relatively small quantities of lime are required, if larger amounts must be applied (1,500 pounds up to one ton or more), calcic limes should be used in order to prevent a surplus content of magnesia in the leaf. The magnesia requirement in the latter case is most readily filled in the fertilizer by inclusion of some magnesian lime or Emjeo.

TABLE 10. CONTENT OF MAGNESIUM OXIDE AND CALCIUM OXIDE IN TOBACCO FROM AIR DRY BASIS, LIME PLOTS. CROP OF 1941.

Liming materials	Grade	CaO	Av.	MgO	Av.
Check	LS	4.96	4.24	1.02	.85
	L	4.16		.84	
	M	4.15		.76	
	LD	3.68		.77	
Hydrated magnesian lime	LS	5.11	4.63	1.94	1.60
	L	4.78		1.59	
	M	4.48		1.56	
	LD	4.13		1.29	
Landplaster—Emjeo	LS	6.00	5.21	1.97	1.75
	L	5.18		1.80	
	M	4.77		1.58	
	LD	4.88		1.64	
Magnesian limestone	LS	5.60	5.07	1.74	1.37
	L	5.06		1.40	
	M	4.96		1.22	
	LD	4.66		1.12	

#### RESIDUAL EFFECT ON STABLE MANURE

In the Tobacco Substation Report for 1940 (Bul. 444:246) the results of the first year of an experiment to measure the residual or after-effect of a preceding ten years of heavy manure application were described. During the first year, although the usual application of commercial fertilizer was used on all the field, those plots which had manure during the previous years gave an increase of 25 percent in yield and 33 percent in grading over the other parts of the field. This experiment is to be continued through a series of years in order to determine how long these effects will last.

The same four manure plots and corresponding controls were measured in 1941, as in 1940. During the summer the differences between the manure plots and the other parts of the field were not so marked as in 1940 but the leaves were a little darker green and the growth a little more luxuriant. When the cured tobacco was on the sorting bench, no very consistent differences between them were observed with respect to color, texture, veins and other quality char-

acteristics. The yields and grading of the plots are shown in Table 11. These data do not show any improvement in grading, but there was an increase of 136 pounds to the acre (7 percent) in yield. This is a smaller increase than was shown the first year but still shows that there was a distinct residual effect two years after any manure was applied.

TABLE 11. YIELD AND GRADING RECORDS OF RESIDUAL MANURE PLOTS AND CONTROLS. CROP OF 1941.

Previous treatment	Plot no.	Yield, lbs. per acre		Percentage of grades								Grade index	
		Plot	Av.	L	M	LS	SS	LD	DS	F	B	Plot	Av.
Manure two years previous	M1	1913		5	4	27	1	39	6	11	7	.386	
	M2	2100		8	4	39	2	35	1	9	2	.462	
	M3	2194	2086	1	2	36	2	38	2	12	7	.381	.410
	M4	2138		3	2	38	1	39	2	9	6	.409	
No previous manure	Mch1	1819		4	2	29	3	40	7	9	6	.384	
	Mch2	1931		8	3	35	3	36	3	10	2	.443	
	Mch3	2063	1950	2	2	33	3	39	5	12	4	.382	.409
	Mch4	1988		8	3	31	2	39	1	12	4	.425	

#### FURTHER EXPERIMENTS ON STARTER SOLUTIONS

From previous trials with starter solutions it was learned that by necessity they should be rather weak (Bul. 444:260-264). Recognizing this fact, in repeating the tests in 1941 the solutions used in the setter barrel were of a weak concentration.

The following solutions and concentrations were employed:

- I. Nitrogen as nitrate of soda, 3 pounds per setter barrel (50 gallons of water)
- II. Nitrogen and potash; 3 pounds nitrate of soda and 1 pound of sulfate of potash per setter barrel.
- III. Nitrogen, phosphoric acid and potash, according to following formula:
 

50	pounds superphosphate (16 percent $P_2O_5$ )
32.5	pounds nitrate of soda
13.5	pounds sulfate of potash
4.0	pounds Emjeo (magnesium sulfate)

100.0

Three and a half pounds of this mixture were used for 50 gallons of water, furnishing plant food at a ratio of 5-10-7.

On a field of rather low fertility, that had been previously fertilized at the rate of 2,500 pounds of 8-4-8 per acre, rows of tobacco were planted, using plain water for check rows and, for alternating rows, the various solutions. Thus, there was one check row, followed by a row of solution I, a row of solution II and a row of solution III. This procedure was repeated so that triplicate test rows were obtained for check and treatments. The field was planted on June 4 and harvested on August 21. Yield and grading records

are given in Table 12. From the data presented it is seen that starter solutions were decidedly beneficial. The results expressed in terms of relative crop values are:

Check = 100

I. Nitrogen = 128.5

II. Nitrogen and Potash = 128.0

III. Nitrogen, phosphoric acid and potash = 127.1

TABLE 12. YIELD AND GRADING RECORDS OF STARTER SOLUTION TESTS. CROP OF 1941.

Treatment	Plot no.	Yield, lbs. per acre		Percentage of grades								Grade index	
		Plot	Av.	L	M	LS	SS	LD	DS	F	B	Plot	Av.
Check	A	1790	1832				13	40	7	20	20	.213	.300
	B	1759				26	3	34	12	14	11	.316	
	C	1947				33	4	48	3	10	2	.372	
N	I A	1862	2016	2	2	32	5	39	7	11	2	.383	.405
	I B	2063		8	7	29	3	42	2	7	9	.448	
	I C	2123				34	3	54	1	8		.385	
N & K	II A	1792	1918			32	6	33	10	10	9	.348	.367
	II B	1892		2	2	25	4	36	11	12	8	.344	
	II C	2070		5	2	31	4	44	2	11	1	.408	
N, P & K	III A	1805	1883			28	6	31	12	11	12	.326	.371
	III B	1816		5		33	5	32	9	9	7	.393	
	III C	2029		4	1	29	6	46	5	9		.395	

The results indicate that using relatively weak starter solutions is beneficial, although it is strongly suggested that nitrate of soda alone will benefit both growth and quality more effectively than mixtures of several plant foods. In using nitrate of soda in the setter barrel it is suggested that a stock solution be made up, such as 50 pounds to 50 gallons of water, of which not more than 3 gallons should be used for a setter barrel.

#### USE OF CARBON BLACK TO ACTIVATE GROWTH EARLY IN THE SEASON

It is commonly known that black surfaces absorb heat rays more efficiently than surfaces of other colors, with a minimum of absorption when the surface is non-colored or white.

Similarly, a dark colored soil absorbs more heat than a light colored soil. If, therefore, a light colored soil could be darkened by some artificial means, it should absorb more of the sun's heat. This would probably benefit plants set out early in the season when cold nights would be a setback to growth. A black material placed around plants would cause the soil to absorb and retain more heat (depending on moisture) and build up a reserve for the night.

In order to determine what effect, if any, a blackened area around tobacco plants might have on growth and quality, a preliminary trial was made in the spring of 1941.

A quite uniform field was selected (field IV) and fertilized with a balanced mixture (2,500 pounds of 8-4-8 per acre). Seven rows of tobacco (150 feet long) were used for the experiment. To four of these carbon (Solium<sup>1</sup>) was distributed in bands 4 inches wide on either side of the row. Alternate rows were left untreated as controls.

The carbon black was applied at the rate of 850 pounds per acre, although it is claimed that a 200-pound rate would be sufficient for other crops. The band on the surface of the soil was undisturbed for about ten days (carbon black applied on June 2 and cultivated the first time on June 12). In the early stages of growth it was observed that at least three of the four treated rows produced somewhat larger plants than the controls. It was thought that the material might have induced more favorable conditions for nitrification. Hence, concentration of soil nitrates was determined at three different times during the growing season. The results are listed below:

Treatment	Pounds per acre of Nitrate Nitrogen			Seasonal average
	June 10	June 17	July 8	
Control .....	53.6	95.8	51.0	67
Carbon black .....	56.4	157.6	85.6	100

This material (Solium), claimed not to contain nitrogen as an impurity, apparently greatly accelerated the processes of nitrification—about 50 percent for the trial period.

The tobacco was harvested on August 18 and, after proper curing, it was sorted in the Station warehouse. As judged on the sorting bench, tobacco from the control rows was considered to be of very good quality, with occasionally a few starved (yellowish) leaves. The quality of tobacco from the carbon black treated rows was considered only fair. It was somewhat coarse, (as contrasted to the desirable smooth, silky texture) and a considerable number of starved leaves was found. The grading and yield data are found below.

Treatment	Yield per acre	Percentage of grades								Index	
		L	M	LS	SS	LD	DS	F	B	grade	crop
Control .....	2172	9	4	31	2	37	1	10	6	.435	944.8
Carbon black..	2311	7	6	28	1	37	1	13	7	.398	919.8

It is shown that the carbon black treatment produced a somewhat higher yield than the control, although the increase is not big enough to cover the additional expense for material and labor. However, if the grading had been proportionally improved, that *per se* might have warranted its use. The grade index for the control was almost 10 percent higher than that of the carbon black. This gives the crop index (product of grade index and yield figure) for the control a value of about 945 as contrasted to that of the treatment, 920. These values, on a relative basis with 100 for the control, allow only 97.4 for the carbon black application.

<sup>1</sup> "Solium" is a commercial grade of carbon black furnished by Binney & Smith Company of New York.

The result of one season's trial suggests no real advantage in employing a material, such as carbon black (Solium), for tobacco. No doubt, however, the material may be used to facilitate biological activities on soils where there is need for such a speeding up and for crops other than tobacco. With respect to tobacco in general and cigar leaf tobacco in particular, quality is often more important than yield.

It is possible that, in the present test, most extensive production of nitrates occurred at a time when the plants were still too young to fully utilize the plant food. This is indicated by the exceptionally high level of nitrates on June 17 (157.6 pounds per acre); and, by the fact that the percentage of "darks" (LD and DS) was no higher for the carbon black tobacco than it was for the control, thus indicating that there was no oversupply of nitrogen when the crop was maturing. (A high percentage of "darks" is usually considered an indication of an oversupply of nitrogen in the soil.) It is also possible that a portion of the extra-early production of nitrates under the carbon black treatment may have been leached below the root zone by June rains.

#### FURTHER INFORMATION ON IRRIGATION OF TOBACCO

In previous reports<sup>1</sup> it has been shown that irrigation of tobacco during dry periods can be quite effective and profitable if measures are taken to prevent depletion of nitrate in the soil during the process. Further data along the same line were obtained by an experiment in 1941, although the season as a whole was not a good one for irrigation because the rainfall was sufficient and well distributed except for one period in mid-July.

On July 23 the tobacco on one coarse sandy field was wilting badly and was therefore used for this test. One-third of the field was irrigated by running water down between the rows (method described in Bul. 391). Another third was not irrigated at all. The last third was irrigated as mentioned but, in addition, nitrate of soda at the rate of 150 pounds to the acre was sprinkled into and dissolved in the water before it sank into the soil. The purpose of this was to replace any nitrate that would be leached away. The rate of water application was equivalent to about one and one-half inches of rainfall. It was planned to repeat this a week later but timely rains made irrigation unnecessary.

When the cured crop was sorted, a high percentage of starved yellow leaves from the plots irrigated without nitrate indicated that the fertilizer had been leached away. Such starvation was not observed on the irrigation plots to which nitrate of soda had been added. The yield and sorting records are shown in Table 13.

<sup>1</sup>Conn. Sta. Bul. 391, p. 67 and Bul. 44, p. 264.



The data show that, on a sandy soil, irrigation alone may be more harmful than beneficial. It resulted in a reduction of 26 percent in the value of the crop (crop index). But when nitrate of soda was added to the water there was an increase of 15 percent in the value of the crop. Both yield and grade index were materially increased. These differences are larger than we have had in tests of previous years but are all in the same direction. It is not likely that such large differences would be found if the soil was heavier and less prone to leaching. However, it is the lighter soils which the grower usually irrigates, because they are the ones that first suffer from drought.

TABLE 13. YIELD AND GRADING RECORDS OF IRRIGATION PLOTS. CROP OF 1941.

Treatment <sup>1</sup>	Yield, pounds per acre	Percentage of grades								Grade index	Crop index
		L	M	LS	SS	LD	DS	F	B		
Not irrigated	1556	3	2	30	1	33	7	12	12	.362	563.27
Irrigated alone	1526	—	—	16	1	39	13	15	16	.273	416.60
Irrigated with addition of nitrate of soda	1649	5	3	32	1	33	5	14	7	.393	648.06

<sup>1</sup>Relative values based on crop index.

Not irrigated	115
Irrigated alone	74
Irrigated nitrate	100

The results of all experiments to date warn against the danger of this type of irrigation on "leachy" soils unless the nitrate content of the soil is kept up.

#### LUMARITH AS A SUBSTITUTE FOR GLASS IN SEED BED SASH

In our report for last year (Bul. 444, p. 269) the advantages of lumarith (a light, transparent, plastic, cellulose acetate) were described. The results of the first year's test were favorable and warranted a continuation of the trial. Sections of the seed bed were therefore covered in 1941 with the same lumarith sash that we had used in 1940. The sash had been kept over winter in a curing shed. By spring the plastic had become very brittle. Unless the sash was handled very carefully, the lumarith tended to split and break loose from the wooden frames. Before the seed bed season was over, most of the material had become unserviceable and had to be replaced with glass sash. Unless this fault can be overcome, lumarith can hardly be recommended because it is obviously impracticable to re-cover the frames each year.

There were no observable differences in growth of the plants in the seed beds covered by lumarith. They did not grow any faster than those under glass, nor was there any other difference in vigor, color of leaves, etc.

It has been claimed by some experimenters that lumarith might have some effects that would be expressed in improved vigor in the field after transplanting. In order to test this, six rows of plants from the lumarith section of the bed were set in a field, each row alternating with a row of plants of the same age from the glass-covered parts of the bed. As far as could be observed there were no differences in growth between the alternate rows in the field during the summer. Each was harvested, cured and sorted separately. The yield and sorting results are shown in Table 14.

TABLE 14. YIELD AND GRADING RECORDS OF TOBACCO FROM LUMARITH AND GLASS COVERED BEDS. CROP OF 1941.

Sash covering	Yield, pounds per A.	Percentage of grades								Grade index	Crop index
		L	M	LS	SS	LD	DS	F	B		
Lumarith	1853	3	2	40	2	39	2	10	2	.421	780.09
Glass	1913	1	1	35	2	39	2	11	9	.398	761.37

The yield was higher in the rows from the glass-covered beds but the grading a little better on the lumarith. The differences are probably too small to be significant or to indicate that any improvement in yield or quality was produced in the crop by growing the plants under lumarith.

## TOBACCO DISEASES IN 1941

P. J. ANDERSON

Diseases of tobacco caused very little damage during 1941. Downy mildew was prevalent for a short time in the seed beds but caused little trouble later and was almost absent from the fields. Not a single case was seen or reported of wildfire, blackfire, frenching, leaf blotch, sore shin or wet stalk rot. Of quite minor importance were damping-off, bed rot, black and brown root rot, hollow stalk and the various forms of pole rot.

### Some Experiments on Control of Downy Mildew

The most common method of applying PDB is to place the crystals in shallow wire baskets, about 2 inches wide and 10 inches long. These are supported by nails driven into the side boards of the beds. Although the early experiments showed that, under favorable temperature conditions, one basket to each two sashes is sufficient when one ounce of crystals to each four square yards of bed is used, many of the growers are using one basket to each sash, "staggering" them on alternate sides of the bed. The degree of control obtained has varied greatly from farm to farm depending on (a) tightness of beds, (b) intervals between application, (c) time of starting treatments, (d) length of treatment, etc.

On the whole, however, it must be admitted that control of mildew by PDB, according to the method of application we have used, has not been entirely satisfactory, especially when the nights are cold. It is possible that a more thorough distribution of the gases in the bed might give better control.

In some tobacco sections of the country where cloth covers are used over tobacco beds, the crystals are distributed over the entire surface of a thin cloth which, in turn, is covered with a heavier cloth. This method gives a wider distribution of the crystals and, possibly, a more nearly equal gas concentration in the air than can be obtained by the basket method. This is reported to give satisfactory control. Since this method would not be very practical under our glass sash, a modification of the same principle was tried in our beds at the Experiment Station in comparison with the wire basket method.

Light frames  $3 \times 5\frac{1}{2}$  feet in size were covered with a thin cheese cloth. These frames were then supported under the glass above the plants by spike nails driven into the sides of the bed. The frames were spaced one to each 6 feet of bed. One ounce of PDB crystals was distributed evenly over each cloth frame every second night. In another section of the bed a similar amount was used in baskets, one to each sash, but staggered. A third section of bed was left untreated as check. Treatments were applied on May 14, 17, 19, 21, 23, 26, 28, 30 and on June 1, 3, 6, 9 and 11. The bed was inoculated thoroughly on May 15 and on June 2 by sprinkling all the bed with a suspension of spores in water.

On May 24 mildew began to appear on the check bed and continued to spread until there was 100 percent infection. On June 1 there was a little infection on the basket section. None appeared on the cloth section.

The cloth method thus gave better results than the basket method in this test, indicating that more thorough distribution of crystals may improve the degree of control.

**Removal of PDB Crystals.** The usual practice is to remove the PDB from the beds about 8 A. M. daily. If the crystals are kept in the beds until it becomes too warm there is danger of plant injury from too great a concentration of the gas. This is indicated when the plants begin to fade to a yellowish cast. But, on cloudy days it does no harm to keep the crystals in the bed all day.

One progressive grower suggested that we should gauge the time of removal by the thermometer rather than the clock. To test his conviction he kept thermometers in the beds and removed the baskets only when the temperature came up to 80° F. He did not have mildew in any of his beds during the setting period although he had mildew there regularly for several preceding years, and there was

mildew in the beds of his neighbors. Unfortunately, he did not leave any untreated beds for a control this year.

Another grower found it too laborious to remove the baskets of crystals each morning and, therefore, he left them in permanently but was careful to raise the sash and ventilate the bed during the day. He escaped mildew. The method is rather wasteful of material but saves labor.

**Liquid Sprays.** Although a considerable list of liquid sprays have been tried here for control of mildew, none of them has given satisfactory control. Most of them have caused leaf burning, or other types of injury, or have failed to control mildew. Nevertheless, new ones have been tried each year. One which we tried this year was the most satisfactory of any of the sprays we have yet found. This is a copper soap spray called "Flordo" which has been used with good results in Florida.<sup>1</sup> In our tests this was prepared as follows: 1 pound of copper sulfate was dissolved in 15 gallons of water, and 4 pounds of soap (soap flakes) in 15 gallons of water. These two solutions were then stirred together. A small amount of curd that was formed was removed by straining through cheese cloth.

This was sprayed on the growing plants twice a week for four weeks. These plants and the unsprayed controls were inoculated twice with spores of mildew. The unsprayed checks finally became 100 percent infected. Although the section sprayed with Flordo did not remain entirely free from disease, infection was light and did not cause serious damage. Some of the leaves were burned by the spray but not severely. Results indicate that this material should be subjected to further trial. A good spray might appeal to some of the growers more than the fumigation method now in general use.

#### Exit Wildfire

This is the first season in 20 years that the writer has failed to see or hear report of a single case of wildfire in the Connecticut Valley. After its spectacular outbreak and destructive spread in the early Twenties, wildfire began to be less prevalent each year. This gradual tapering off process continued until during the late Thirties it was unusual to find more than a dozen cases any year, but each year there were always a few. Now it seems to have dropped out completely. Such behavior for any plant disease is quite unusual in the annals of plant pathology.

#### DELAYED FROST INJURY TO TRANSPLANTS IN JUNE

During the first two weeks of June light frosts were reported from scattered localities after the plants were started in the fields. Frost was most general on June 12, but there were also reports of

<sup>1</sup>Shippy, W. B. Flordo Spray, Fla. Agr. Exp. Sta. Press Bulletin 504 (1937) and 547 (1940).

quite localized frosts on other nights. The chill was not severe enough to cause the leaves to turn black and die. In fact, no one reported injury at that time. But within the next two weeks growers began to bring in plants with peculiar markings which they were unable to diagnose. Having forgotten about the earlier frosts, they feared that another new disease had appeared here.

The most characteristic symptom of frost injury is the presence of broad white chlorotic bands, equally distributed on each side of the midrib and usually confined to the basal part of the leaf (See Figure 2.) The tips and upper parts of the leaves are commonly green and normal. The outer margin of the basal part may also be green, but frequently the chlorotic bands involve all the web of the basal part. The whitened, affected base fails to expand as the leaf



Figure 2. Delayed frost injury showing at center of young tobacco transplant.

grows and, therefore, appears narrow and pinched, making a sort of a handle to a spoon-shaped leaf. The margin of the injured part is usually wavy and the leaf variously distorted. The older, bottom leaves may be quite normal. Only the leaves that were quite young

and tender at the date of the frost are involved. The affected area of the leaf may be solid white or may be variously speckled and peppered with green.

In general the injury was found to be confined to limited parts of the field. All three types of tobacco were affected. Observations later in the year showed that all leaves above the chlorotic lower ones developed normally, and the plant as a whole was not seriously injured.

A similar condition occurred here in 1926 following a heavy frost on June 16 (See Tobacco Sta. Bul. 8:55 for description and illustration).

## MOSAIC OR CALICO DISEASE OF TOBACCO

The season of 1941 was marked by unusual freedom from damaging diseases of the tobacco crop. But, contrary to this general statement, mosaic (known also among growers as calico, mongrel, brindle, gray top and rust) was more abundant than we have known it to be for many years. Its sudden prevalence aroused renewed interest among growers, and the Substation received numerous inquiries about its cause, spread and prevention. In view of this interest and the fact that we have been experimenting for several years on a method of control, about which information has not been previously published, this seems an opportune time to present an up-to-date picture of our present knowledge of tobacco mosaic.

It has been 12 years since we published a comprehensive article on mosaic (*Tobacco Sta. Bul.* 10, p. 75). In the interim the problems of mosaic diseases of plants have been extensively investigated by scientists. Their findings have greatly advanced our understanding and changed our conceptions of the causes of these diseases. Numerous articles by plant pathologists, physiologists and chemists have been published in journals and bulletins, but they are not available to many of our readers or are too technical for the practical tobacco man. The purpose of this article is to present in non-technical language the latest pertinent facts about mosaic.

**Symptoms.** Every experienced tobacco grower recognizes an ordinary calico plant in his field. The mottled appearance of the leaves, with splotched areas of greenish yellow between other areas of normal green, is familiar to all (Figure 3, top). Symptoms show all gradations of intensity. Sometimes the differences in shade of color are barely noticeable; at the other extreme, the light chlorotic areas of the leaf may be bright yellow to almost white and therefore stand out very sharply in contrast. Also, the pattern or mosaic, formed by the irregular alternating chlorotic and normal parts, is so extremely varied as to defy description. In some types, however, the darker areas have a tendency to follow the veins (Figure 3, bottom).

Attempts have been made to distinguish and name several types of mosaic according to differences in leaf symptoms, such as pattern or color of the chlorotic areas. There are, however, so many intergradations between field types that, except for the mosaic specialist, such a classification adds nothing to clarity of understanding. In severe cases young leaves may become distorted or "puffed" and are inclined to be abnormally narrow. Plants affected early become dwarfed, do not make a normal growth and are practically worthless when mature. Late infestations with only a few of the top leaves showing the symptoms, frequently not severe, are commonly known as "gray top" by growers.



Figure 3. Tobacco leaves showing patterns of mosaic: Top, the most common pattern of mottling; center, red rust, a symptom of calico marked by irregular, dead, brown patches; bottom, the pattern in which the dark areas follow the veins.



But there are other symptoms which growers do not always realize are manifestations of the same disease. This is particularly true of the necrotic or dead spots of various shapes and colors that develop in the leaf tissue. These are of two general types and have been called white rust and red rust. The white rust, also called fleck, is marked by numerous small white spots peppered over the leaf. In the red rust type the spots are larger, more irregular and of a reddish brown color (Figure 3, center). This may be so severe as to cause death of the whole leaf. The red rust type, if more diffused, is sometimes called mosaic burn and appears often on the upper leaves of the plant. Burning of the top leaves usually results from the spread of mosaic while topping. Since such leaves often show no mosaic mottling, the grower is not aware of the connection.

**Extent of Damage.** Although mosaic recurs here every year, it is rarely a disease of major importance. In this section as a whole only a fraction of one percent of all the plants set contract the disease at a sufficiently early stage to suffer serious deterioration in quality or yield. An occasional field is found where the loss is heavy, and it is the occurrence of such cases that gives importance to the disease and the need of control measures.

The extent of damage to a plant depends largely on the stage of the plant's development at the time it becomes infected with the disease. A plant infected in the seed bed probably never attains sufficiently normal growth for its leaves to be suitable for cigar wrappers or binders. They might possibly serve as cheap scrap chewing tobacco. On the other hand, if infection occurs at the time of topping, the damage may be practically negligible, though, even at this late stage, considerable "burn" of the top leaves may result.

There are three sources of loss. (A) Dwarfing of the plants reduces yield. (B) The dead spots (white and red rust) reduce the value of the leaves somewhat in proportion to prominence of the spots. (C) A mottled leaf, when cured, has uneven colors, dead finish and is not elastic.

In a three-year experiment in Maryland, McMurtrey<sup>1</sup> found that early infestations caused a loss of 55 percent of the gross value of the crop. In another three-year test in Kentucky, Johnson and Val-leau<sup>2</sup> found that early infection reduced the yield 31 percent and the total value 40 percent. Even when infection occurred at topping time, the quality of the tobacco was reduced 25 percent. The losses in both of these tests are calculated on the basis of 100 percent infection — a condition that rarely occurs naturally — but they serve to

<sup>1</sup> McMurtrey, J. E. Effect of mosaic disease on yield and quality of tobacco. Jour. Agr. Res. 38:257-267.

<sup>2</sup> Johnson, E. M. and W. D. Valleau. Effect of tobacco mosaic on yield and quality of dark fire cured tobacco. Ky. Agr. Exp. Sta. Bul. 415. 1941.

show that any degree of infection is damaging both to yield and quality.

**The Causal Agent.** Ever since tobacco mosaic was described 60 years ago, a considerable variety of theories on its cause have been advanced and supporting evidence published to back each theory. Only in the last few years has the true cause been demonstrated and all the other alleged causes relegated to interesting past history. Mosaic is now known to be caused by a virus, appearing in the plant sap as minute rod-like particles of a protein nature which have the remarkable characteristic of being able to multiply indefinitely in the living cells of all parts of the tobacco plant.

According to recent investigations<sup>1</sup> an increase in these particles is at the expense of chromoprotein, which is in the protoplasm of all the cells of the plant. Whether or not the virus is composed of living organisms, it has definite parasitic properties. The cell cannot synthesize the virus *de novo*; a small particle of it must be introduced into a living cell in order to cause infection. Here it multiplies and spreads to other cells. When introduced into a tobacco plant, the virus spreads rapidly to the roots and to the newly developing leaves, but extremely slowly to leaves already fully grown. The symptoms described above will appear on the leaves that were growing rapidly (rust) and on leaves which were very small (mottled) when infection occurred. All new growth will likewise be mottled. The particles of virus in the plant juice, which are designated as protein molecules by some investigators, are very minute, so small that they will pass through the pores of very fine filters. It would require about 125,000 of them placed end to end to span one inch.

**Spread and Overwintering.** Mosaic is extremely infectious and will spread from the most minute quantity of the virus transferred to the cell of a growing plant. From this point it quickly spreads in the sap to the roots and to the growing point of the plant. Once a plant is infected, it never recovers, since it has no way of eliminating the virus. When, in handling infected plants, the juice gets on the hands of the workman, he need only touch or rub his fingers over the leaf of a healthy plant to infect it. Any virus introduced into seed beds would be thoroughly disseminated by weeding or pulling the plants for setting. When there are a few infected plants established in the field, the virus may be spread around on hands, clothing, tools, etc. during any cultural operations. Fortunately, it is not spread by many of the common disseminating agents of fungous diseases, such as wind and splashing rain. There is no conclusive evidence that it is spread by insects.

Although the methods of dissemination in beds and field are well known, the means by which mosaic first gets its start each spring are

<sup>1</sup>Woods, Mark W. and H. G. DuBuy. Synthesis of tobacco mosaic virus protein in relation to leaf chromoprotein and cell metabolism. *Phytopath.* 34:978-996. 1944.

not so obvious. Its method of overwintering, however, gives some clue. In cured or dried leaves the virus will remain infective almost indefinitely. Thus Johnson and Valleau<sup>1</sup> found it is still active in tobacco that had been kept in a case for 52 years. Its longevity is probably the most important fact to be kept in mind in connection with the prevention of mosaic. It will even survive in manufactured tobacco, such as cigars, cigarettes and chewing tobacco, although in smaller amounts than in the barn-cured tobacco.

Valleau has shown that in Kentucky the use of "natural" chewing tobacco by workmen tending the beds is the common source of initial bed infection. Badly infected beds have been observed in Connecticut where growers have used tobacco refuse, raked from the floor of curing sheds, as fertilizer. We visited a field in the summer of 1941 where one part of the field had 75 percent mosaic plants, while other parts of the field were relatively free of the disease. The infection was traced to the fact that the owner had ground up tobacco stalks from the previous year's crop and applied them as a fertilizer to the area of lowest fertility at time of setting the plants. This insured a thorough distribution of mosaic. In some sections of the country perennial weeds belonging to the tobacco family are sometimes infected, and the virus is transferred to tobacco plants while weeding. This does not appear to be a common source of infection here, however.

A great deal of investigation has centered on the ability of the virus to live in the soil over winter. It has been shown that in many soils the virus survives the winter, but probably only a very low percentage of infection may be traced to soil-borne virus from a preceding mosaic crop. The virus may be in the still undecomposed residues of the plants or may have been washed out into the soil itself. The percentage of virus that winters over in this way is obviously quite small. Otherwise, it would not be profitable to grow tobacco year after year in the same field. Every grower has seen fields where infection was about 100 percent on the suckers that grew up after a crop was harvested. Yet, a crop planted on that same field the next year would as likely as not be free from mosaic. If all the virus had remained active in the soil, there would be few healthy plants in the new crop.

No doubt, the greater percentage of virus becomes inactivated in the soil, especially in the upper layer which is stirred and well aerated before setting. Virus, like other protein substances, is decomposed by the action of soil organisms and thereafter cannot cause infection. The residues of tobacco plants quickly decompose when incorporated in the soil, especially if the soil is moist and well aerated. Investigators have shown that the type of the soil may have an influence on the ability of the virus to live over. This may explain the report

<sup>1</sup>Johnson, E. M. and W. D. Valleau. Mosaic from tobacco one to fifty-two years old. Ky. Agr. Exp. Sta. Bul. 361:264-271. 1935.

by many growers that certain spots in their fields are prone to show mosaic more or less every year. The soil in these areas may be more suitable for keeping the virus in an active condition.

**Control.** No spray or dust is effective in controlling mosaic. Control measures must start with an understanding of the nature of the disease and, particularly, its method of dissemination as described above. Methods should aim, first, at preventing the introduction of the disease into the bed or field in the spring and, secondly, at removing infected plants which may spread it from plant to plant in the field.

The following preventive measures have been found useful and are offered as suggestions to growers whose crops are troubled with mosaic.

1. Tobacco refuse from curing sheds, sorting rooms or other sources should be kept entirely away from the seed beds. Sometimes tools or bed sashes, which have been stored in sheds without being carefully cleaned, may carry bits of diseased leaves to the beds.

2. If the stalks of a preceding crop are to be used on tobacco land for fertilizer, they should be plowed under the surface as thoroughly and early as possible to give them every opportunity to decay. If stalks are from a mosaic crop, or if results from this practice in previous years have been bad, it is better to use them on land intended for other crops.

3. Men who are weeding, pulling or setting, or handling the crop in any way, should be discouraged from using tobacco. This applies to the use of the leaf in any form, but particularly in chewing and pipe tobacco.

4. Plants should never be set from a bed known to contain the disease. Mosaic plants are difficult to detect in the seed bed. If only a few are found, it is quite certain that there are many other diseased ones which have not been "spotted." Rather than risk spreading the disease to the field it is better to sacrifice the whole bed. Second and third pullings from a bed are much more likely to have mosaic than the first ones, because of spread while pulling.

5. Just after the plants start in the field, "rogue" out any diseased ones. About once a week, while the plants are small,<sup>1</sup> a workman should systematically inspect each row, pull up every calico plant and put it in a basket. This should be done before cultivating or performing other operations which might spread the disease. The person who rogues should avoid touching any healthy plant, otherwise he may spread more disease than he removes. The basket of diseased plants should be dumped far from any growing

<sup>1</sup> If plants are large there is too much danger of spread by carrying them through the field.

tobacco. The workman should thoroughly wash his hands before handling healthy plants again.

We regard this "rogueing" as a very essential measure of control. Both the Experiment Station farm and commercial growers have used it with success for many years. If it is carefully done, no other control measure should be necessary the remainder of the season. Rogueing is practicable only where the percentage of infected plants is small, but this is the condition in at least 95 percent of Connecticut fields. Where the percentage is very high, it is better to plow under the crop and replant from healthy beds.

6. If rogueing has not been successful, avoid spreading the virus from diseased to healthy plants during field operations, even at the time of topping and suckering. If there are several men working, one of them may handle all the diseased plants ahead of the others and not touch the healthy ones. This measure is of less importance than the preceding since the direct damage at this stage is not great, but it has some bearing on the practice mentioned in the next paragraph.

7. Every precaution should be taken to keep the entire tobacco farm free from mosaic. By carrying out sanitation practices the grower reduces chances of infection from refuse, which holds no danger if leaves and stalks are disease-free. Moreover, a soil which does not harbor mosaic cannot spread infection. Some Connecticut tobacco farms have had practically no mosaic for years because the owners will not tolerate any sources of infection.

**Mosaic Resistant Varieties.** The perfect method of controlling any disease is to find or develop a strain or variety of the crop which is naturally immune or so highly resistant to the disease that need of further control measures is permanently eliminated. A resistant strain of tobacco must also have satisfactory commercial qualifications. The incorporation of good characters in a resistant strain may be the most difficult part of a breeding program.

Until 1933, it was currently believed that all varieties of tobacco were equally susceptible to mosaic and therefore offered little chance for the development of a resistant strain. Then Nolla and Roque<sup>1</sup> found in Columbia, S. A., a variety of tobacco called Ambalema. This is so highly resistant to mosaic that common symptoms are absent, except for an occasional indistinct mottling of the lower leaves. Although Ambalema does not have qualities comparable to those of any types grown commercially in the United States, its factor of resistance may be bred into our present varieties by hybridization. Several experiment stations and the United States Department of Agriculture are endeavoring to get suitable commercial resistant strains by using Ambalema as a parent. This hybridization pro-

<sup>1</sup> Nolla, J. A. B. and Roque, Arturo. A variety of tobacco resistant to ordinary tobacco mosaic. Jour. Dept. Agr. Puerto Rico 17:301-303. 1933.

gram requires years, selection and, although good progress has been made, we do not know of any investigator who is yet ready to distribute seed of a resistant variety which he considers to have suitable commercial qualities.

At this Station we are attempting to breed a resistant Broadleaf strain. Starting with a cross between Ambalema and John Williams Broadleaf, we have back crossed three times on Broadleaf. Many selections are satisfactorily resistant but they have not been tested long enough to make sure that they possess proper commercial qualities.

### INSECT INVESTIGATIONS DURING 1941

A. W. MORRILL, JR., AND D. S. LACROIX<sup>1</sup>

The cooperative investigations on the control of insect pests of tobacco in the Connecticut River Valley, which were commenced in 1936 by the Connecticut Experiment Station and the Bureau of Entomology and Plant Quarantine of the United States Department of Agriculture, were continued during 1941.

Various dust mixtures were tested on shade-grown and open field tobacco in an effort to obtain a simultaneous control of both potato flea beetle, *Epitrix cucumeris* (Harr.), and tobacco thrips, *Frankliniella fusca* (Hinds). Because of the impossibility of applying sprays effectively to tobacco late in the season without causing mechanical injury to the plants, spray materials were not continued in the program. Naphthalene and carbon disulfide were tested as soil fumigants for the control of the eastern field wireworm, *Limonius agonus* (Say). In addition, the field surveys and general observations of insect conditions on tobacco were continued, as in previous years.

#### Flea Beetle and Thrips Experiment

Treatments were tested in an attempt to determine a single control measure for use against both the potato flea beetle and tobacco thrips.

Although setting of tobacco plants was commended as early as May 15 this season, winds of 15 to 26 miles per hour blew almost constantly, preventing dusting until June 12. Thereafter, the treatments were applied at semi-weekly intervals when the weather permitted until July 11, when heavy showers prevented further dusting operations. Harvesting of the leaves was completed shortly thereafter. A total of eight applications was made with each insecticide. The rate of application of dusts commenced at 6 pounds per acre at the beginning of the season and increased to 12 pounds per acre as the plants grew. Applications were made by means of rotary

<sup>1</sup>Bureau of Entomology and Plant Quarantine and Connecticut Agricultural Experiment Station, respectively.

hand-operated dusters in the early morning, usually before 9 A. M. Standard time, because this period of the day is freest from disturbing wind movements.

Each treatment, as in all similar tests conducted since 1937, was replicated six times on plots arranged in a Latin square. Each plot consisted of one bent of shade-grown tobacco, or one-fortieth acre. In conformity with the methods used in previous experiments, four primings of three leaves each were harvested from each of 10 plants, also chosen at random, in every plot. These leaves were then cured, fermented and processed in the usual commercial fashion. They were later examined for evidences of insect injury, which may be either more or less apparent on the cured leaf than on the green.

Three criteria were used in judging the effectiveness of the treatments, as follows: (A) the populations of flea beetles and thrips as sampled at intervals during the growing period of the crop, (B) the percentage of cured leaves showing evidence of feeding injury by these insects, and (C) the percentage of the total potential wrappers (four per leaf) showing injury severe enough to damage the value of the wrappers. Counts of live insects were made on (10) sample plants chosen at random in each plot immediately before and, again, 24 hours after each application of the insecticides. The percentage of leaves injured by feeding was determined by examining all leaves harvested from 10 plants per plot, as previously mentioned, after curing and processing were completed. In the case of the last-mentioned criterion, as applied to flea beetle feeding, injury severe enough to have caused commercial damage when properly located on the leaf was counted as actual damage when it was not so located. Thus this criterion is a measurement of the true severity of the injury, but overstates the commercial importance of such injury in the case of the flea beetle.

The use of pyrethrum dust gave better control of thrips in terms of wrappers damaged than did any of the other treatments, except that it was not significantly better than the treatment in which pyrethrum was alternated with cubé. There was no significant difference in the effectiveness of these two treatments and that of the dust mixture containing 1 percent of rotenone in controlling the flea beetles. All treatments were better than no treatment.

#### Control Experiments on Open Field Tobacco

A somewhat similar experiment on the control of tobacco thrips and the potato flea beetle was conducted on plots of open field tobacco in Buckland, Conn. Each treatment was replicated four times on plots one-fortieth acre in size that were arranged in two randomized blocks. The treatments used consisted of semi-weekly applications of the dusts shown in Table 1 during the period August 1 to August 16. These dusts were applied by means of rotary hand-operated dusters

at rates of 3 to 30 pounds per acre, depending upon plant size. The effectiveness of treatments was determined by counts of living flea beetles and thrips on 10 plants per plot just before and 24 hours after each application. The results from this experiment are given in Table 15. These data show a trend somewhat similar to that of the results on shade-grown tobacco.

The large decrease in thrips population after the fourth application was probably due to heavy thunder storms which came in cloudburst proportions, not only washing off many of the thrips but also spattering the tobacco foliage with mud to such an extent that it was difficult to make accurate counts.

The figures accumulated in 1941 on Broadleaf tobacco bear out the same facts as were found in 1940 on Havana Seed tobacco, i. e., pyrethrum does serve to hold thrips in check to a certain degree, but is not entirely satisfactory.

Pyrethrum dust with a mineral diluent left a white residue on the tobacco leaves. This had a tendency to wash down towards the base of the leaves during rains and accumulated there. During dry weather, the white residue was plainly visible over all the foliage.

#### Wireworm Control Experiments

On June 6, twelve plots, one-fortieth acre in area (33 x 33 feet) were treated with naphthalene at the rate of 800 pounds per acre, applied in furrows 9 inches deep, approximately 9 inches apart, and later harrowed. Twelve others were plowed and harrowed, but were not treated with naphthalene. Twelve similar plots were treated with dichloroethyl ether, applied in the setting water at the rate of 100 cc. of ether per gallon, about one-third to one-half pint of the mixture being applied around each plant. This ether was made more soluble in the water by the addition of a commercial cat-ion-active emulsifying agent at the rate of 3 parts by volume of the agent to 100 parts of water. Twelve additional plots were treated with setting water containing 2 grams of para-Aminoacetanilide and 12 were plowed, harrowed, and set with plain water. The 48 plots were arranged in 12 randomized blocks. The effectiveness of the treatment was determined by counting the wireworms obtained in sifting the soil from six units of 1 cubic foot each, located at random in the central portion of each plot. The plots were set with tobacco three days after treatment with naphthalene, and no difference could be noted in the vigor of the plants on the treated and untreated plots. However, the ether caused considerable burning of the roots and resetting was necessary.

While air temperatures were unseasonably high from early spring, soil temperatures remained below 70° until the day of setting, after which they rose to between 70° and 75° for some days. On June 18 and July 9 the average reductions in wireworm infestation were de-



TABLE 15. EFFECT ON OPEN FIELD TOBACCO OF INSECTICIDAL TREATMENTS FOR THE CONTROL OF THE POTATO FLEA BEETLE AND TOBACCO THRIPS, BUCKLAND, CONN., 1941.

Treatment	Number of living insects on ten plants for indicated application														Post-treatment total <sup>1</sup>		
	No. 1		No. 2		No. 3		No. 4		No. 5		No. 6		No. 7				
	Before	After	Before	After	Before	After	Before	After	Before	After	Before	After	Before	After			
FLEA BEETLE																	
Pyrethrum dust <sup>2</sup> with:																	
Clay base	6	3	1	3	0	0	2	18	25	3	123	21	21	3	223		
Exhausted pyrethrum flowers base	3	2	1	1	1	0	3	10	14	0	105	11	8	4	160		
Denicotinized tobacco-dust base	7	1	1	0	1	2	2	7	42	3	121	42	37	1	260		
Undusted check	3	5	0	1	0	2	6	2	50	25	94	50	73	31	339		
THRIPS																	
Pyrethrum dust with:																	
Clay base	18	3	34	19	28	22	129	9	3	1	106	39	71	30	494		
Exhausted pyrethrum flowers base	18	4	28	28	27	29	80	7	3	1	115	25	55	51	453		
Denicotinized tobacco-dust base	14	9	33	31	26	28	129	7	2	1	101	63	57	24	511		
Undusted check	15	8	24	54	54	65	127	18	9	2	96	76	78	96	707		

<sup>1</sup> Does not include the first count.<sup>2</sup> 0.5% pyrethrins I and II in all tests

terminated by soil sampling. The results of the samplings taken on June 18 were not significant. Those of the sampling taken July 9 are shown in Figure 4. While the reduction in wireworm infestation was barely significantly greater in the plots treated with naphthalene than in those not treated and somewhat more so in the plots treated with the ether, the reductions obtained were not sufficient to be of much practical value to the grower.

In September a new set of 36 plots was arranged in randomized blocks in a field which had been heavily infested at setting time and in which 6 resettlings were required to get a stand. Twelve plots were treated with carbon disulfide immediately following the removal of the tobacco plants after harvest. Twelve plots were plowed to a depth of about 1 foot and 12 were left untreated, except that all 36 were harrowed before and after the plowing. Temperatures remained fairly high during the period, the average daily departure being 2.7 Fahrenheit degrees above normal. A reduction in the wireworm population of 87 percent was obtained in this experiment, which is considered to be highly significant (Figure 4).




Figure 4. Results of treatments applied for the control of wireworms in tobacco fields, Buckland, Conn., 1941.

#### Insect Abundance During 1941

A survey to determine the severity of damage by insects to open field tobacco in the Connecticut River Valley was conducted during August, following the method used in previous years. This survey

showed that the damage by insect pests was less severe than in 1940 and much less severe than for a number of years preceding. As in 1940, many pests were late in becoming established and some of the usual pests were not present in any appreciable numbers.

Percentages of leaves showing injury and of those showing commercial damage caused by the insects of major importance were as follows: Flea beetles, 20 percent and 7.97 percent, respectively; thrips, 6.54 percent and 1.84 percent; grasshoppers, 7.31 percent and 1.01 percent; and hornworms, 2.66 percent and 0.59 percent.

The Japanese beetle, *Popillia japonica* (Newm), again was observed feeding on tobacco, but not in such large numbers as in 1939 and 1940. As in previous instances, injury was confined to the tops of the plant and was very moderate.

The wireworm *Limonius agonus* (Say), which has been often reported under the name *Limonius ectypus* (Say), is frequently a severe pest of newly set tobacco in the Connecticut River Valley. In 1941 this insect was an especially serious problem, probably because of the unusual earliness of the setting of the young plants in the field. Many fields were attacked so heavily that restocking was necessary five and six times and, since the resetting must be done by hand rather than by the usual setting machinery, these operations were unduly expensive. Attack was usually confined to small areas in certain fields, as is normally the case but, in some instances, whole fields of fifteen to thirty acres were affected. This latter condition was seen especially in fields which have been set alternately to both potatoes and tobacco. Damage by the seed corn maggot, *Hylemya cilicrura* (Rond.), was not reported or observed, nor was there any damage by crane fly larvae or "leather-jackets," which in this region usually are the larvae of *Nephrotoma ferruginea* (F.), *sodalis* (Loew), or *Tipula georgiana* (Alex.).


The potato flea beetle, *Epitrix cucumeris* (Harr.), and the tobacco thrips, *Frankliniella fusca* (Hinds), were very scarce following heavy rains early in the season, but became almost normally abundant just before harvest. Injury by the flea beetle was approximately as general as last year, but leaf infestation was not as heavy. On the other hand injury by thrips was unexpectedly severe on cured leaves. Whereas excessive feeding by this insect will cause the leaf to become thin and papery and thus of little value, even moderate injury causes a white streaking of the leaf. When populations are small, this streaking is confined to the midrib section and is of little importance. However, during dry seasons, if no control measures are applied, the thrips may crowd farther out onto the leaf, and injury is caused to the wrapper portion of the leaf. This occurred during the latter part of the 1941 season.

Grasshoppers were rare until near harvest time, when more feeding than normally was observed in outside rows, probably ow-


ing to the drought conditions. The injury in these fields was caused mainly by the red-legged grasshopper, *Melanoplus femur-rubrum* (De G.) and, rarely, by the Carolina grasshopper, *Dissosteira carolina* (L.). Cutworms in seedbeds were not reported and few cases of injury by these insects to the growing plants were observed.


Other usual pests of tobacco in this region, not reported in 1941, include the tarnished plant bug, *Lygus pratensis oblineatus* (Say), the garden springtail, *Bourletiella hortensis* (Fitch), and the stalk borer, *Papaipema nebris nitela* (Guen). The spined stink bug, *Euschistus variolarius* (Beauv), the spotted cucumber beetle, *Diabrotica duodecimpunctata* (F), and the potato aphid, *Macrosiphum (Illinoia) solanifolii* (Ashm.), were found doing only very slight damage.

No specimens of the tobacco budworm, *Heliothis virescens* (F.), were observed. A few light infestations of the tobacco hornworm, *Protoparce quinquemaculata* (Haw.), were observed on sun-grown tobacco before harvest, and several were found on the later suckers. Although unlikely, it is possible for these worms to overwinter in this region and to emerge as moths during succeeding years. Such infestations on suckers are, therefore, potential sources of damage to subsequent tobacco crops, and the suckers should be carefully destroyed when they appear after harvest. At the same time, growers should take care not to destroy larvae found crawling in curing sheds when these larvae are covered with the small, white cocoons of the parasite, *Apanteles congregatus* (Say). Contrary to the general conception, these cocoons are not hornworm "eggs" but a potential source of a hornworm control of considerable value and effectiveness in this region.













University of  
Connecticut  
Libraries

---



39153029345131

